

Paper March 10

1844

W. S. H

Dean

An inquiry  
 Into the Medical virtues of the  
 Gal. Vinchennae and Sulphur Quinque.  
 In the case of Intermittent  
 And Remittent Fevers.  
 Submitted as an  
 Inaugural Thesis  
 to the examination of the  
 Reverend Frederick Penrose, Rector.  
 The Trustees,  
 And Medical Professors of the University of  
 Pennsylvania.  
 On the first of November 1844.  
 In the Degree of Doctor of Medicine.  
 By Charles Henry Rich  
 of  
 Philadelphia.





To William Fisher M.D. of the University of Pennsylvania.

Dear Sir,

Prompted by inclination, and impressed by the highest respect for Your professional talents, I claim as Your pupil, the privilege of dedicating to You my Inaugural Dissertation. But, independent of all other considerations, I am impelled to inscribe this essay to You by gratitude for the many opportunities of improvement, and the unceasing attention which You have so kindly afforded me during the term of my Medical studies under Your direction. I am conscious that it is so unworthy Your acceptance, that in its faults as You have been accustomed to do in those of its author, and I shall have consolation in reflecting, that Your candour to excuse, is no less eminent than Your abilities to judge.

That you  
and I  
that the  
nation  
you

The  
C  
H  
S  
a  
W

That you may long continue to enjoy health,  
and the honors awarded to your merit, and  
that the happiness of Yourself and Family may  
continue to increase, is the unfeigned wish of  
Your much obliged friend and grateful

Pupil  
Charles C. Mohr.

To  
The Reverend Philip V. Meyer D. D.

This Essay

Is respectfully inscribed.

As a grateful testimony of the  
Affection and Esteem.

With which his uniform Friendship

And politeness

Has inspired

The Author.

The  
month  
to his  
Matron  
The cha  
tiz an  
Genera  
Ca  
An  
Sta  
St

Botanical arrangements.

The genus *Cinchona* belongs to Linnaeus' four-  
teenth Natural Order *Centurata*.

In his sexual system he places it in the class  
*pentandria*, and Order *Monogynia*.

The characters which he gives of the fructifica-  
tion are as follows:

General Characters.

*Calyx*.—Superior, one-lobed, five-toothed,  
permanent.

*Corolla*.—Monopetalous, tubular, funnel-  
shaped, deeply five-lobed, tube long,  
obtusely angular, segments lance-  
olate or linear.

*Stamens*.—Filaments five in the middle of  
the tube, anthers elongated, wider  
shorter than the tube, or projecting  
beyond it.

*Pistil*.—Germ inferior, top-shaped, obscure.

*[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]*

by ang

Puisay

Epim

Co

Co

Ho

Cap

ly angular, style the length of the stamens, stigma  
thick, either bifid or entire.

*Pericarpium*. Capsule crowned with the calyx,  
two-valved, two-celled, valves open-  
ing at their inner side, turned in  
at their edges, and separating when  
ripe, so as to have the appearance  
of two capsules.

*Seeds*. Numerous in each cell, oblong, com-  
pressed, biconvex, attached to a cen-  
tral oblong receptacle.

#### *Essential Characters.*

*Calyx*. Top-shaped, five-toothed.

*Corolla*. Tubular, five-lobed.

*Stamens*. Five, inserted into the middle of the  
tube.

*Capsule*. Oblong, two-valved, two-celled, many-  
seeded.

*[Faint, illegible handwriting on lined paper, likely bleed-through from the reverse side.]*

*[Faint, illegible handwriting on lined paper, likely bleed-through from the reverse side.]*



Natural History

This tree is very lofty, sending off large branches covered with rough brown bark: the leaves vary from an ovate to an elliptical shape, the larger approaching more to the former, and the smaller more to the latter figure. They are all entire, serrated, smooth on the upper side, on the under, tomentose, and stand in pairs upon petioles. The flowers are produced in panicles, and stand upon slender pedicels. The calyx is small, bell-shaped, and cut at the margin into five minute segments: the corolla is funnel-shaped, consisting of a long cylindrical tube, divided at the limb into five segments, about an ovate, or oblong spreading, on the upper side, on the under, woolly, and fringed at the edges. The filaments are bristly, placed in the middle of the tube, and furnished with oblong anthers, twisted in a spiral manner.

the  
what  
with  
sails,  
alled,  
It is a  
to a t  
broth  
to, rec  
Mr. M  
but is  
and h  
erend  
intoms  
cullie  
the in lo  
see, jee  
Having  
obtaind,

the germen is water: the style is foliiform, some  
what longer than the stamina, and furnished  
with a round stigma, the capsule divides into two  
parts, the cells of which are separated by a par-  
allel partition. the seeds are small and numerous.  
It is a native of peru, growing most abundantly  
on a long chain of mountains, extending to the  
North and South of Lima, where its trunk frequent-  
ly reaches in bulk the body of a man. According to  
Mr. Bird, the soil in which these trees thrive  
but is generally a red clayey, or rocky ground,  
and especially on the banks of small rivers de-  
scending from high mountains. This author also  
informs us that the most proper season for  
cutting the bark is from September to December,  
the only time in the whole year of some intermis-  
sion from the rain in the mountains.  
Having discovered a spot where the tree was most  
abundant, they first build huts for the workmen,

and  
hall  
they led  
having  
she then  
then or  
a farm  
the rain  
with to  
provide  
a large  
fifty per  
the one  
the hall  
ground  
yard less  
with sh  
the steps  
hall as

and then a large hut, wherein they put the  
bark in order to preserve it from the wet, but  
they let it lie there as short a time as possible,  
having first cut a road from the place where  
the trees grow, through the woods, sometimes  
three or four leagues, to the nearest plantation  
or farm house, in the low country, whither, if  
the rain permits them, they carry the bark fresh  
with to dry. These preparations being made, they  
provide each Indian, they bring the cutters, with  
a large knife, and a bag that will hold about  
fifty pounds of green bark: every two Indians  
take one tree, whence they cut, or skin down  
the bark as far as they can reach from the  
ground; they then take sticks about half a  
yard long each, which they tie to the tree  
with strong withs at proper distances, like  
the steps of a ladder, always slicing off the  
bark as far as they can reach, before they go a

new st  
dian lo  
they de  
the bag  
of this  
the muc  
the bas  
per, it  
low-oc  
colue  
time in  
the sam  
ting, she  
he carry  
dying  
the Gre  
On the  
they see  
the whic

new step, and thus mount to the top, the Indian below gathering what she other cuts, this they do by turns, and go from tree to tree till the bag is full, which, when they have plenty of trees is generally a days work for one Indian. As much care as possible must be taken that the bark is not cut wet; should it so happen, it is to be carried directly down to the low country to dry, for otherwise it loses its colour, turns black, and rots; and if it lie any time in the tree without being spread, it runs the same risk. As soon while the Indians are cutting, the mules, if the weather permits, ought to be carrying it down to the place appointed for drying it, which is done by spreading it in the open air, and frequently turning it. On the trees being entirely stripped of their bark they soon perish, and as the number of these trees to which access could be had, was said to be

not a  
a sup  
mand  
heroi  
by lo  
from  
to be  
agate,  
uable  
We se  
what  
effia  
so wi  
Some  
ous, w  
she of  
nab  
or she  
she ju



not very considerable, it has been supposed that a sufficient quantity of bark to supply the demand, could not long be procured. Leonardine, however asserts that the young trees do not die by losing their bark, but send out fresh shoots from the base, and as those which are supposed to become old have time to disseminate & propagate, we trust, the fear of exhausting this valuable medicine is wholly unfounded.

We seem to have no satisfactory account at what time, and by what means the medical efficacy of the Peruvian bark, which is now so well established was first discovered.

Some contend that its use in Intermitting Fevers, was known to the Americans long before the Spaniards possessed Peru, but that they communicated this knowledge from the Europeans, and on the contrary it is asserted by others, that the Peruvians never supposed it to be fit for

ang  
quan  
ding  
is off  
ding  
me th  
time  
Sutton  
he cur  
shun  
me lo  
time b  
into a  
broome  
Thim  
size  
ing as  
to drink  
the after

any medical use, but thought, that the large quantities exported thence, was for the purpose of dying, and they actually made some trials of its effects in this way. Bonnamy says, that according to an ancient tradition, the Americans owe the discovery of this remedy to the lions, which, some naturalists pretend, are subject to a kind of Intermittent fever, of which they were observed to be cured by instinctively eating the bark of the cinchona. But Desfontaines states that the use of the bark was learned from the following circumstance. — Some cinchona trees being thrown by the winds into a pool of water, lay there, till the water became so bitter that many <sup>of</sup> refused to drink it. However, one of the neighbouring inhabitants being seized with a violent paroxysm of fever, and finding no other water to quench his thirst, was forced to drink this, by which he was perfectly cured. He afterwards related the circumstance to others.

qua p  
q p  
il p  
has a  
the p  
pamia  
bische  
have w  
bemit  
china,  
Quinqu  
tributa  
vite, in  
regulate  
into L  
Hobbs p  
as, beca  
this the  
the first

and prevailed on some of his friends who were ill  
of fevers, to make use of the same remedy, in which  
it proved equally successful. The use of this well-  
known remedy was, however, little known, till about  
the year 1655, when a signal cure having been per-  
formed on the Spanish Prince's lady, the Countess de  
Briehen, at Lima, it came into general use, and  
hence was distinguished by the appellation *pulvis*  
*beniéfico*, also called *Antea China china*, a *China*  
*china*, *Mina mina*, a *Minamina*, and *Quina quina* or  
*Quinquina*. On the recovery of the Countess, she dis-  
tributed a large quantity of the bark to the Ser-  
vants, in whose hands it acquired a still greater  
renovation, and by them was first introduced  
into Europe, and thence called *pulvis Peruvianus*,  
*pulvis Patruum*, and also *Cardinal de Lugo's pow-  
der*, because that charitable prelate bought a large  
quantity the fourteenth, when Dauphin, was said to be one  
of the first in Europe who experienced its efficacy.

quantity  
as the  
the bar  
some  
plots  
in par  
a yellow  
out her  
town  
better  
avote  
the lot  
ish  
where  
as it  
water  
is  
ble  
any

quantity as a great resource for the religious poor  
at Rome.

This bark is brought to us in pieces of different sizes,  
some rolled up in short short quills, and others  
flat: the outside is brownish, and generally covered  
in part with a whitish mofe: the inner side is of  
a yellowish, redish, or rusty iron colour. The best  
cut breaks close & smooth, and pieces friable be-  
tween the teeth: the inferior kinds appear when  
broken of a woody texture, and in chewing, of sep-  
arate into fibres. The former pulverises more easily than  
the latter, and looks, when powdered, of a light brow-  
ish colour, resembling that of cinnamon, a some-  
what paler. It has a slight smell, approaching  
as it were to mustiness, yet so much of the aro-  
matic tincture as not to be disagreeable. Its taste  
is considerably bitter, astringent, and very disa-  
greeable in the mouth, and accompanied with some  
degree of aromatic warmth, yet not sufficient to

person  
shot of  
we can  
there, is  
shot of  
shot of  
St. Louis  
practice  
much  
most  
to  
gather  
like the  
by comp  
gen, for  
and of  
it, more  
my little  
is more  
Louis's



proves its being ungrateful. Besides this bark, that of several other species of *cinchona* have been recommended for medical use by different authors, especially the latter *peruviana* Aublet, also that of the *cinchona baribaca* a Samarra bark, that of the *cinchona Humboldtii* produced at St. Lucia, and that of two or three other species produced at Santa Fe. The first of these is in much larger and thicker pieces than the common, most of them are coarse, though not called together like the quilled bark. They break short, like the best common bark, and appear evidently composed of three layers, the outer is thin, rough, frequently covered with a mossy substance, and of a reddish brown colour. The middle is thicker, more compact, and of a darker hue, it is very bitter and resinous. The innermost layer is more woody and fibrous, and of a lighter red. Louis' *Matéria Médica*, p. 275.

the p  
some  
ious  
the sub  
the m  
ting. The  
presul  
but m  
is inter  
devote  
is in a  
infusion  
the ad  
tincture  
stringe  
the exte  
of the p  
this is on  
the, and

In preparing this bark, the middle layer, which seems to contain the greatest proportion of resinous matter, does not break so readily as the rest; a circumstance to be attended to, lest the most active part should be left out in sifting. The sea bark to the taste discovers all the peculiar flavour of the common Peruvian bark; but much stronger. An infusion in cold water is intensely bitter, more so than the strongest decoction of the official bark. Its astringency is in an equal degree greater than that of the infusion of the common bark, as is shown by the addition of the Sulphur Ferri. The spirituous tincture of the sea bark, is also proportionally stronger than that of the pale. The quantity of matter extracted by rectified spirits from the powder of the former, was to that of the latter, as three to two in one experiment, and as 129 to 130 in another, and yet, on infusing the two residuums of

the p  
on h  
which  
that y  
mid of  
Bupiole  
aspect  
brill p  
men, in  
found  
and den  
ing to  
obscure  
by being  
useful  
ter, for  
at peace  
was a  
12, 1  
1811. 1811

the first experiment in boiling water, that of the  
red bark gave a liquor considerably bitter, and  
which struck a black with Sulphuric Acid; while  
that yielded by the other was nearly tasteless, and  
void of astringency.

Respecting the medicinal properties, we have several  
respectable authorities, showing, that as the red  
bark possesses the same properties with the com-  
mon, in a much higher degree, so it has been  
found of more efficacy in the cure of Intermittents,  
and hence it is thought to be that, which accord-  
ing to Aret. the Spaniards called *Cascarilla*  
*colorada*, and was probably the wine original-  
ly brought to Europe, and which proved so suc-  
cessful in the hands of Sydenham, Astruc & Les-  
ter; for it appears from the testimony of the old-  
est practitioners, that the bark first employed  
was of a much deeper colour than the common.  
\*Lew. Drugs & Matters upon. †Med. Trans. v. 20. p. 161.

the sea  
shot of  
time given  
the Doctor  
in the  
just, he  
ingly go  
about a  
as the go  
endingly  
bring the  
in the sea  
ually the  
the latter  
sing given  
of the sole  
himself  
which is  
where is

The sea bast was first imagined by Saunders to be  
shot of the full grown trees, the branches and young  
trees yielding the summer bast, but this opinion  
the Doctor seems afterwards to have abandoned, for  
in the third edition of his pamphlet on this sub-  
ject, he says: that he had lately seen some exceed-  
ingly good sea bast, imported by a Spanish mer-  
chant, a considerable portion of which was as small  
as the guillica bast in common use. It was ex-  
ceedingly resinous, and gave evident proofs of its  
being the quilt of the larger sea bast which was  
in the same chest. — If the pale and sea bast were  
really the produce of the same species of birch, the  
latter differing from the former only by acquir-  
ing greater maturity, we should find the disposition  
of the colour of the pale bast to correspond propor-  
tionally with its thickness, in the side of the quilt,  
which is certainly not the case. The birchona tre-  
chana is described and figured by Saquin, &

Distin

the sta-

tion of

the Phil

ture su

histori

to its

satiation

you ca

how the

is apper

also see

The perso

hiding

shifting

forme a

by thea

taini a

will in



Greatly bright, it grows in Samaria, where it is called  
the Star-like bark. The Chinese *Huichunda*, a bark  
tree of *St. Lucie*, a figure of which may be found in  
the Philosophical Transactions, also in *Visier's* observa-  
tions sur la Physique, affords a bark which is  
likewise said to have been used with advantage.  
In its second state it is considerably emetic and  
cathartic, properties which it retains in some de-  
gree on being dried, so that the stomach does not  
bear this bark in large doses, and in small ones,  
its effects are not such as to give it any particu-  
lar recommendation.

The Persian bark yields its virtues both to  
boiling and cold water, but the decoction is  
shittier, gives out its taste more readily, and  
forms an ink with a chalybeate more success-  
fully than the fresh cold infusion. This infusion con-  
tains at least as much extractive matter, but  
more in a state of solution, and its colour ex-

the  
the q  
an of  
sunder  
an for  
Mila  
abstract  
the al  
of the  
a few  
in the  
is per  
to the  
then sh  
mild  
water,  
del of  
tum of  
Bartle

standing with a chalybate becomes darker, while  
that of the decoction becomes more pale. When they  
are of a certain age, the addition of a chalybate  
renders them green, and when this is the case, they  
are found to be in a state of fermentation & efflu-  
escence. If a caustic alkali, or lime, precipitate the  
extractive matter, which, in the case of the caustic  
alkali is dissolved by the further addition  
of the alkali. Lime water precipitates less from  
a fresh infusion than from a fresh decoction, and  
in the precipitate of this last, some mild earth  
is perceptible. The infusion is by age reduced  
to the same state with the fresh decoction, and  
then they deposit nearly an equal quantity of  
mild earth and extractive matter, so that lime  
water, as well as chalybeates may be used as a  
test of the relative strength and purifiable na-  
ture of the different preparations of the different  
salts. Accordingly, cold infusions are found

by express  
refusals  
of the po  
by length  
with the  
Some wat  
ter, appe  
offic em  
para the  
bar sup  
a bill  
of 25 offic  
astings  
of as ma  
ua of bo  
from mas  
vian bas  
which go  
Edinburgh

by experiments to be less perishable than decoctions,  
infusions and decoctions of the *ra. baist.* than those  
of the pulp, those of the *ra. baist.* however are found  
by length of time to separate more *mila* earth  
with the *lime* water, and more *extracted* matter.  
*lime* water, as precipitating the extractive mat-  
ter, appears an equally improper and disagre-  
able menstruum. Water has been found to sus-  
pend the resin by much less gum than has  
been supposed. Rectified spirit of wine extracts  
a bitterness, but no astringency, from a residuum  
of 20 affusions of cold water, and water extracts  
astringency, but no bitterness, from the residuum  
of as many affusions of rectified spirit. The resi-  
ua of both are insipida.

From many ingenious experiments made on the *Pru-*  
*rian* *baist.* by Dr. Boing, published in a dissertation  
which gained the prize-moat, given by the *Bar-*  
*Edinburgh* *Acad.* Dispensatory.

since  
difficult  
ascertain  
for his  
to some  
in the  
Gale  
Monday  
Gale  
Antip  
some to  
the anti  
the, and  
change  
unf  
some ex  
ing, yet  
and its  
it that

seian Society of Edinburgh is 791. the following  
different menstrua upon the Peruvian Bark is  
ascertained with greater accuracy than had be-  
fore been done, and it appears, that with respect  
to comparative power, the following fluids act  
in the order in which they are placed.

Eulogia spirit of Nitric. - Caustic liq. French  
Brandy. - Rhineish Wine. - Vinegar diluted. -

Eulogia spirit of Nitric. - Mild volatile alkali.

Rhetical spirit of wine. - Mild vegetable alkali.

Some water. -

The anti-scorbic powers of vinegar and base uni-  
ted, are double their own taken separately  
though the bark on its first appearance was and  
was for some time afterwards, was reported by  
some eminent physicians to be a dangerous reme-  
dy, yet these prejudices are entirely gone away,  
and its character is now universally established  
so that the disputes which now subsist are

10/10/19  
 was in  
 visit  
 I am  
 prepared  
 the new  
 settled  
 in the  
 The me  
 schemes  
 infusing  
 and co  
 hours,  
 10/10/19  
 because  
 This me  
 the why  
 enough  
 intention



applied to the mode of preparation, and the manner in which it is most advantageously administered.

I am now to speak particularly of two recent preparations from the cinchona, whose powers in the cure of intermittent fevers have been greatly extolled. The first to which I allude was called in the shops, *sal. cinchonae*.

The method employed by a manufacturing chemist of this city for its preparation, consists in infusing a certain quantity of the *Cortex Peruviana* in a given quantity of water, for 24 hours, straining the liquor thus obtained, and evaporating to dryness, which leaves a castaneous residuum, being the article in question. This medicine has been extensively employed by the physicians of this city. In a time it was thought to be very efficient, and indeed in the extensive practice of a gentleman of the profession

of my  
is al  
Best  
as im  
half of  
for an  
by ne  
employ  
transac  
has a  
solution  
for she  
aspetate  
in a  
prostit  
tivity  
a p  
subj  
has all

of my acquaintance, in the Northern Liberties,  
is almost superseded the exhibition of the  
Bast in substance. On its first introduction it  
was employed in minute doses. The fourth, to the  
half of a grain were said to be proper proportions  
for an adult. Finding by repeated trials that she  
by no means answering the expectations of those  
employing it, the doses were from time to time  
increased, until they were finally established at  
ten or twelve grains every two hours, in full re-  
solution.

In the course of the summer of 1792, the great  
expectations which had been formed of it, abated  
in a considerable degree in the estimation of those  
practitioners whose opinions I have had an oppor-  
tunity of consulting. I have not had the advantage  
of personal experience with the medicine, but shall  
submit what information on the subject I have  
been able to collect from the practice of others.

Order  
obliging  
expected  
partial  
me to  
the  
operation  
had good  
to be  
survived  
from the  
of things  
all other  
the opinion  
view of  
Happened  
to and  
in order  
to a

Doctor George Wither, of the Boston Liberator, has  
obligingly communicated to me the result of his  
experience with the article, after a fair and im-  
partial trial, which he has politely permitted  
me to transcribe. — I have employed the Sal Bi-  
chuene in a great number of cases, with its  
operation in a great number of instances I have  
had great reason to be pleased, but particu-  
larly so when the basis in substance, or more bulky  
preparations could not be retained on the stomach,  
from the irritability of that organ. Where this state  
of things existed, I have found it to remain, when  
all other preparations were rejected by vomiting. —  
The opinion of Doctor Alexander Wright, first physi-  
cian of this city, concurs exactly with the preceding.  
He gave a mixture of Sal Bichuena  $\frac{ss}$  Esti. Ess. Ferrous  
ii. and q. Doctor Zipp, in one particular instance,  
in doses of a dessert spoon full every three hours,  
to a child three years old. —

The first  
Book  
was of  
the  
Spice  
with  
m. - I  
was la  
at an  
ed to  
na, an  
sible  
was tel  
the mo  
admine  
The an  
ter, wit  
stomach  
any at

The following communication<sup>2</sup> from my friend  
Doct<sup>r</sup> S. A. Uhler in answer to a desire I ex-  
pressed of having an abstract of his experience with  
the *Pal. Cinchonae*. — According to your request,  
I give you this brief statement of my experience  
with the *Pal. Cinchonae*. I was called to see  
M. — S. —, who at the time of my visiting her  
was laboring under Remittent Fever. I prescrib-  
ed an emetic &c. and during the apyrexia attempt-  
ed to give an infusion of the Barks, *Serpenta-*  
*ria*, and Orange peel, (which I generally found to  
sit well on the stomach,) but in this instance it  
was totally unable to bear the smallest quantity of  
the medicine. The same difficulty occurred in the  
administration of some other preparations.  
The case eventually took on a typhoid charac-  
ter, with a total prostration of strength, the  
stomach still rejecting every thing, and resisting  
every article usually employed in such cases.

to qu  
l am  
midic  
the pa  
in this

ly wh  
how a  
vins of  
how to  
tea to  
of sube  
I am  
the Volp  
from the  
store.  
mistaken



to quice its irritability. - In this dilemma,  
I commenced the use of the Sal. binchonae: the  
medicine was retained without difficulty, and  
the patient gradually recovered. The form I used  
in this case was as follows:

R. Sal. binchonae ʒij

Mucil. S. Arabic. ʒj

Aqua Mentis. ʒssij. ʒo. It.

Of which a table-spoon full was given every  
hour and a half. - I have not used the med-  
icine often, but from the little experience I have  
had with it, it appears to be pretty well adap-  
ted to cases of the above character, though, to be  
of service it must be given in large doses. -  
I am enabled to speak in more decided terms of  
the Sulphas Quininae, another late preparation  
from the Peruvian Bark, than in the preceding in-  
stance. The prevalence of Intermitents and Re-  
mittent Fevers in the Northern suburbs of the city

his son  
personal  
provision  
a young  
join the  
midwint  
furnish  
Vol. 6  
Digne  
Kant, to  
provision  
Sulphur  
base of the  
barbarous  
the principal  
sublime  
set with  
ing agitar  
plus, sa

this season, afforded me a good opportunity of personally testing this medicine.

Prior to, however, to transcribing my own diary, or giving the experience of others, I will subjoin the formula for the preparation of the medicine employed by the same chemist who furnished me that for the formation of the Val. Linchona. It is as follows:

Digee for 24 hours, a quantity of Peruvian Bark, with four times its weight of water, previously mixed with a sufficiency of Sulphuric Acid to saturate the soluble base of the bark. Strain off the liquor, and add Carbonat. Calcis to neutralize the acid. Filter the precipitate, wash, dry by a moderate heat, & pulverize it: put the powder into a close vessel with six times its weight of alcohol; having agitation it will, by heat gradually applied, raise it to a boiling temperature.

This off  
not and  
and sat  
note and  
Sulphur  
on inter  
without  
it is us  
of nomi  
shitar  
in gett  
knives  
be grade  
age, ha  
The m  
one oad  
O. P. de  
using the  
have been

This effected, remove it from the fire, suffer it to cool and the precipitate to fall down: strain and saturate with sulphuric acid, evaporate and crystallize. By this process the Sulphate of Quinine is afforded, white, and of an intense and permanently bitter taste, and without odour.

It is used in pill and solution: the latter mode of administration is particularly suited to children, where difficulty is often experienced in getting them to swallow pills, and also possesses the advantage of admitting the dose to be graduated with more accuracy to suit the age, habit &c.—

The medium proportion for adults is one, to one and a half grain every hour.

Dr. R. M., in speaking of the article says: "after using the medicine in a great number of cases, I have been highly gratified with the operation of

for  
tunes  
proce  
have  
one  
have  
I have  
Dr. S.  
my son  
by the  
more  
by me  
our la  
nile of  
no.  
was la  
the use  
unbin  
of "polar

the remedy, and cannot speak of it in higher  
terms than by saying, considering the great  
power it possesses over the epidemic, and its more  
than common prevalence during the summer  
and autumn, I do not know what we should  
have done without it.

I have also to acknowledge my obligation to  
Dr. S. — for the following paper, accompany-  
ing his opinion on the Pal. Lincolnae.

Of the Sulphate of Quinine I can speak from  
more extensive experience, but having been ve-  
ry much engaged during the prevalence of  
our last epidemic, I did not make any min-  
utes of cases, but can give you a few from mem-  
ory. — I was called to see Mrs. S. —, who  
was labouring under intermissions prior to the  
usual vaccination. I prescribed Quina in  
combination with Sassafras, Super-tartrate  
of potash, Sulphate of Alumine & Symplicia. Mes-

shots

but

of this

standing

without

and

Palmer

how

was

men

the

of

parry

of

but

your

above

many

you



chata, which I have always found an excellent combination. She continued in the use of this prescription for upwards of two weeks, during which time she was faster in labour, without any beneficial effect. She grew worse, and I resorted in place of the preceding the Pulchate of Livine in doses of one grain every hour during the apyrexia. She took ten pills, & was cured, having had but one chill after the commencement of this use. About two weeks after, she suffered a relapse in consequence of exposure, & gave the Quinine, and she had not another paroxysm.

I could go on to give you cases without number, but do not consider it necessary to burden your paper with a formal history of each, the above case being much the general result. In many cases there was not a repetition of the paroxysm when Quinine and rarely when Potash.

Ap  
and  
had  
and  
I have  
practi  
encom  
astoi  
qu sho  
For sho  
appet  
wine  
the pla  
Sept  
with 2  
bod lo  
wished  
suing  
atima

Upon the whole, I deem it the most important  
and valuable remedy employed in Syphilis-  
tine form, and speak thus highly from a vast  
array of experience with it.

I have consulted the views of several other  
practitioners, who agree with those of I have  
enumerated, but to transcribe them more in  
detail would swell my paper to a size lar-  
ger than is desirable.

For the following patients, in whom I had an  
opportunity of personally testing the me-  
dicine I am also indebted to Dr. S. M., one of  
the physicians of the Quakers of the Don-  
September 24th. Hannah A. — aged 49 years,  
with Luetidarian Syphilis. At this time she  
had laboured 16 weeks under the disease. She had  
received no medical treatment previous to my  
seeing her. The symptoms being such as usually  
attend the disease, I think the enumeration of

of the  
provi  
a succe  
a gnia  
25th No  
probatio  
discon  
rising a  
26th. The  
yesterday  
though  
27th and  
October 3  
plaine  
from the  
we sudje  
the med  
region  
by here

of them unimportant. Finding it indicated, I prescribed a cathartic of Calomel & Rhubarb. I advised her the Pulphas Divinae in doses of half a grain every hour during the operation.

25th. On visiting her to-day, I found that the evacuation had occurred at the usual hour, the disease having undergone no change since the preceding day.

26th. My patient experienced no prurigen since yesterday.

Though I visited my patient daily on the 27th, 28th, 29th, and 30th yet the attack was not unusual.

October 1st. She has not had a return of her complaint since I last saw her.

From the exposure to which persons in her condition are subjected, I thought it prudent to continue the medicine until the 1st, recommending it to be given at longer intervals, say half a grain every hour and a half on the 27th, every two hours

in the  
in day 8  
in the  
then 11  
now  
cited  
25th. 17  
the same  
26th. 17  
the same  
27th. 17  
28th. 2  
even  
October  
To them  
the case  
before the  
Court

on the 25th, every three hours on the 27th, and so on.  
To-day I discharged her from further treatment, cured.  
In the same family with the preceding case were  
three other children, labouring under the same  
disease, whose treatment was similar to their  
sister.

23th. They had an accession of the paroxysm at  
the usual hour.

26th. One of the children escaped a renewal of  
the exacerbation to-day. The third had suffered its  
accession, but in mild form.

27th. One of them had a return of the fever.

28th. 29th. 30th. On these three days they experi-  
enced no symptom of a renewal of their complaint.

October 1st. They continue free from the disease.

In them I also directed that the medicine, as in  
the case of their sister should not be given up  
before the 1st of October.

Discharged from further treatment, cured.

Septemb  
is in the  
of the D  
He had  
previous  
of Linn  
from 2  
much in  
that she  
had 1/2  
to be occ  
16th she  
as attac  
the 27th  
town of  
October 4  
and that  
Henry o  
pland



September 25th Tues. M. —, aged about 30. This was,  
as in the former instances a case of Intermittents  
of the Quinarian form, which had existed for 8 weeks.  
She had been under medical treatment three days  
previous to this date, and had taken the Sympate  
of Quinine, the dose prescribed for her was one  
grain every hour. She stated to me that she was  
much improved since using the medicine, and  
that the paroxysms now occurred later than they  
had before taking the pills. I recommended them  
to be continued.

26th. She informed me that she had suffered  
no attack since yesterday.

The 27th, 28th, 29th & 30th brought with them no re-  
turn of fever.

October 1st. I called again to see her, and unan-  
dered that she had felt no interruption in her cus-  
tomary employments from a return of her com-  
plaint since last I saw her.

The  
greatest  
Septem  
were a  
one to  
them, a  
region  
the other  
the aia  
with the  
had her  
27th Sep  
with the  
from to  
27th Sep  
with the  
their con  
October  
the said

The medicine was continued to this date gradually  
diminished. — Discharge cured.

September 25th. The two children of Mrs. N. —  
were attacked with the epidemic about the  
same time with their mother, the youngest of  
whom, a boy of 4 years of age, had not another pa-  
roxysm after one day's employment of the medicine!  
The other, a girl of 9 years still laboured under  
the disease.

26th the boy remained free from fever his sister  
had her accustomed paroxysm.

27th Neither of them experienced its return.

28th The girl complained of a slight chill and  
fever to-day.

29th They suffered no exacerbation.

30th They continued exempt from a return of  
their complaint.

October 1st On enquiring of Mrs. N. — to-day,  
she said that the children had retained their

health  
The  
sua.  
Septemb  
the  
Antic  
previ  
mical  
in hea  
propert  
eth &  
polar  
zph. the  
was left  
and the  
ed by  
and the  
do about  
the other

health since my last visit.

The caution to prevent a relapse having been observed, I discharged them, cured.

September 25<sup>th</sup> Mrs. E. — and two children.

There were cases of the prevailing epidemic, of the putrid form, which had subsisted two weeks previous to their having made application for medical advice. I directed the Sulphate of Quinine in hourly doses of one grain for the Mother, and proportionate ones for the children.

26<sup>th</sup> I could perceive but little variation in my patients.

27<sup>th</sup> Mrs. E. — states that her last paroxysm was less distressing than any former one. The distress which had somewhat improved, and the youngest experienced no exacerbation.

28<sup>th</sup> The disorder continued to decline in Mrs. E. — she still remained much the same as yesterday, the other suffered no increased attack.

27th.

On the

the diso

October 2

had new

Diehard

Septembe

age, in

plained

sullens

the 27th.

pasture

that 1000

about 1000

and the

the 28th.

some

October 2

one since

29th. They all escaped an exacerbation.

On the 30th they still remained exempted from the disease.

October 5th I paid a visit to this family, and found that none of them had relapsed since the 30th.

Discharge.

September 26th. Mrs. L., a person of 77 years of age, residing in the family of Mrs. C., complained of great pain in the head, back & limbs, dullness and drowsiness. Visiting my patients on the 27th. she informed me that soon after my departure yesterday, a regular paroxysm of Intermittent fever had come on. Immediately after its abating, she took the Quinine pills, and continued their use through the day.

The 28th, 29th & 30th brought no return of the paroxysms.

October 3rd. No relapse having occurred in this case since my last visit, I discharge her cured.

Hydrogen  
which is  
produced  
in the  
of sugar  
days  
tea into  
sustitute  
sustituted  
having

One is  
the only  
both  
that  
situation



September 29th I called to see Mrs. Eliza D.,  
who I found labouring under remittent fever, with  
nausea, pain in the head & extremities. I learned  
on enquiry that this had been originally a case  
of regular quotidian intermittent, which, after 10  
days continuance under that form, had degenera-  
ted into what I now found it. I prescribed an  
emetic, and directed the Sulphate of Quinine in the  
remissions of the fever. I also recommended the fol-  
lowing formula during the exacerbations:

℞ Pulv. Strich. potass. ℥ij  
Rad. Spicac. ʒssj  
Opii. ʒss

Diss in charact. Bro of — one woman to be ta-  
ken every two hours.

Being my patient, she stated to me  
that after the operation of the emetic she felt  
relieved of the fullness of stomach & pains, with  
a returning appetite: the remission was also

unsider  
and a  
October  
visit.  
with a  
and the  
Disab  
the freig  
inter. I  
as they  
ending.  
It may  
not col  
recurren  
a subje  
hair ex  
are acc  
that the  
the mag

unsensibly longer than after her former paroxysms  
and support less from the violence of the fever.

October 1st. No exacerbation occurred since my last  
visit.

15th. I found my patient fast recovering her strength,  
and no return of her disease.

Discharged from further treatment.

The foregoing cases are a few of which I kept min-  
utes. I could have narrated several others, but  
as they offer very little variation from the pre-  
ceding, I have not recorded them.

It may not be amiss to state that these were  
not selected cases, but consigned to me as their  
recommendations were presented.

A subject of vital importance to practitioners, as  
their credit, and the lives of those under their  
care are jeopardised by its desecrated notice. It is,  
that the high price at which this medicine is  
sold, may, and already has caused the adultera-

die

Sept 18

Page 6

ANAL. Calcd for  $C_{10}H_{10}O$ : C, 88.10%; H, 7.41%. Found: C, 88.1%; H, 7.4%.

PLATE 13

8. 10. 1947

1818

of the case

Christ

The Time

3/11/42

1898

us. Not

800

the man

Emice.

see 600

22. 11. 1881

Dec 20

tion of it. In two instances of which I have been  
 apprised by a gentleman of the faculty, which  
 came under his own cognizance, (being present  
 under his medical attendance) the most serious  
 malignancy had little to have resulted from the  
 administration of arsenious acid with the article.  
 In looking over the last number of the Journal  
 of Medical Science, I read the following remarks, which  
 I think not irrelevant to the subject I have been  
 treating:

Messrs. Pelletier & Bouteau have long been con-  
 vinced that a great number of vegetable substan-  
 ces, possessing strong powers over the animal econ-  
 omy, owed their energy to a salifiable base, as  
 the morphine in opium, the strychnine in mus-  
 semine, the veratrine in heliotrop. — They there-  
 fore began to experiment on Cinchona, with  
 the view of discovering a salifiable base if any.  
 They soon found that the cinchonine, a chrys-

tulle  
stence  
they  
acuto  
of its  
a flac  
only  
this  
ing to  
the  
netom  
and so  
and  
and  
matter  
of signs  
If we  
we shall

tollizing principle of Garnil of Lisbon, was a substance of this kind united with an oily matter. They combined this substance with different acids, ascertaining the form, proportions and properties of its salts, which all had a bitter, sub-acid taste, is flavour, proper to the pale bark, which was only used by Garnil. The binchonine itself has this peculiar taste, but much less marked, owing to its high degree of solubility. Pursuing the examination of the pale bark, Messrs L. P. & C. determined also the nature of the other component substances, which, joined to the binchonine are as follows: 1, binchonine united to Nitric acid: — 2, a green oily matter: — 3, binchoninic acid: — 4, Not soluble colouring matter (stannic), 5, Yellowing matter: — 6, Silicate of lime: — 7, Gum: — 8, Starch: — 9, Lignous matter: — 10, ...  
If we enter into a consideration of these substances, we shall see the greater number, viz the binchonine

not, she  
the one  
radical  
not the  
indivisible  
in the  
very same  
principle,  
which was  
the basis  
of every  
theory of  
liberty.  
The alcohol  
contains it.  
Every thing  
principles  
nearly all



sed, the oily matter, the gum, the ligneous mat-  
ter and the starch cannot possess any of the  
medical virtues of the bark, of which they have  
not the taste: the greater number of them are con-  
insoluble. The soluble colouring matter or the tan-  
nin cannot either be regarded as the active substance  
in the cinchona, for if the latter owed its virtues to  
any tannin matter, other barks, very rich in this  
principle, ought also to have the properties of the  
cinchona in proportionate energy according to  
the tanning matter they contain, but which is  
entirely to medical purposes. There remains the  
Stearate of Lime, but this salt when pure has no  
bitter taste, and is besides insoluble in alcohol.  
The alcoholic tinctures of bark, therefore, do not  
contain it, and they are necessarily febrifuge.  
Every thing induces us to believe that the active  
principle of the bark is the cinchonine; conse-  
quently it is to this substance & to its salts, that

the allusion  
that is made  
to the "Jews"  
yellow hair  
table, and  
and other  
capacity for  
and its  
convinced  
The analysis  
lost the  
Greece, and  
India by  
last is the  
Drops and  
Nepatra  
more, and  
stained.

the attention of the physician and his therapeutic researches ought to be directed. Phila<sup>a</sup> Iowa, p. 263  
Muss<sup>g</sup> Pelletier and Baccator, in analyzing the yellow bark, obtained a substance not crystallizable, and differing from cinchonine in physical and chemical properties. It was a salifiable base, its capacity for acids being different from cinchonine, and its salt more bitter. This substance has been denominated Ninine.

The analysis of red bark presented an extraordinary fact—the simultaneous presence of cinchonine & Ninine, and each in a greater quantity than of-  
funda by the pale and yellow bark. The red bark is therefore justly regarded the best.

Process employed by Mr. Pelletier for procuring the cinchonine and Ninine.

Repeated alcoholic tinctures of the bark are first made, and by evaporation, the alcoholic extract is obtained. It is in this extract that the cinchonine

in the  
sion be  
suitab  
quanti  
drops of  
ing is  
with a  
few mi  
one age  
fills a  
toxin  
in solu  
it with  
and trac  
the solu  
the a lea  
purity.  
is some  
miller

as the *Stimula* is found, which exists in the *Prun-*  
*ian* bast. To obtain the alkaline substance of a  
suitable purity, the extract is boiled in a certain  
quantity of water, to which has been added a few  
drops of hydrochloric acid: the liquor after cool-  
ing is filtered—then concentrated, and treated  
with an excess of magnesia—boiling it for a  
few minutes, the liquor is again suffered to cool,  
and again filtered. The precipitate received on the  
filter is composed of *Stimula*, calcined magnesia,  
tannin and lincolnic acid. Wash the precipitate  
in cold water, dry it in a sand bath, and treat  
it with boiling alcohol, which dissolves the alkali,  
and leaves the magnesia and tannin united to  
the coloring matter. It remains but to evaporate  
the alcohol, and obtain the *Stimula* of a superior  
purity. The alkali of the *Lincolnic* thus prepared  
is sometimes impure from an admixture of oily  
matter—to separate which, and purify it, see

most  
with  
progress  
to the  
more, the  
stock  
are, tri  
the July  
great as  
with  
the admi  
ning and  
the fence  
national  
the ch  
the fine  
Asian  
the asse  
water.

must dissolve it anew in an acid largely diluted  
with water, and treat it for the last time with  
magnesia and alcohol as before mentioned.

In the same work from which these extracts are  
made, may be found another mode of preparing the  
article as directed by M. Badolier in the *an. de che-  
mie*. vol. *XV*. p. 278.

The sulphate of Stront has now been employed by a  
great number of French physicians in fevers of type,  
with considerable uniformity of success. M. Doublet  
has administered the salt in doses of two grains mor-  
ning and evening, which have been sufficient to cause  
the fever to cease. To prevent a relapse, however, it was  
continued for some days longer.

In other experiments made at the Hospital de la Char-  
ité, five grains have been given daily in simple quin-  
tidian fevers, and continued for 5 days. Even the very  
first dose the recurrence of the paroxysms was pre-  
vented.

the Royal a  
 of the Sulp  
 but the g  
 ynnia, we  
 to autumn  
 large doc  
 was seen  
 to all the  
 work is sa  
 th. judom  
 as agree  
 the disc ma  
 to be use  
 more of the  
 is division  
 shows a  
 in perpa  
 & Rural.




Doctor Bally has reported nine cases of fever to the Royal Academy of Medicine, cured radically by the Sulphate of Quinine. It appeared to Dr. Bally, that ten grains given in five doses during the apyrexia, were sufficient to check at once the fever. In actual intermittent, however, he considers that larger doses will be necessary. Two great and obvious advantages of the salt of base over the powder are, the smallness of the dose, whereby the stomach is saved from nausea and oppression, and the freedom from astringency, it being rather tonic of an aperient than of an astringent nature. The dose may be carried to six or seven grains in the 24 hours, which would be equal to about three ounces of the powder. In pernicious fevers, when it is advisable, during a short remission, to introduce a large quantity of the Cinchona, this new preparation will be of incalculable value! Dr. Duval, second naval physician at Brest, has

not of  
 but for  
 of living  
 and the  
 which  
 would not  
 she all the  
 in the  
 serious  
 the the  
 a person  
 three-  
 how. So  
 any and  
 man that  
 in. The  
 informa  
 has pro  
 me wh

sent up a memoir of several cases of Intermittent  
fever of different types, cured by the Sulphate  
of Quinine. The medicine was prepared by Messrs. Basse  
and Solomb, of Brest, who employed a new process,  
which consisted in boiling the yellow bark in water  
acidulated with acetic acid, and precipitating  
the alkaline principle by ammoniacal St. Remondin,  
St. Galli, and St. Robiquet have also furnished  
various notices of this interesting discovery.

That the sulphate Quininae is a valuable remedy,  
a perusal of the authorities I have given, and the  
above-enumerated cases will be sufficient to  
show. In its employment I refrained from using  
any auxiliary medicines, save in one instance, in  
order that the result might be the more conclu-  
sive. True it is that several practitioners have  
informed me that they have seen instances when  
it has produced diarrhoea, and some other symp-  
toms which the bark is known to give rise to



in some cases, but those instances are of rare oc-  
currence, I have never seen any such effects arise  
out of its use. —

Dr. Gold

by

James Henry Freese

Charleston, S.C.

